WE CLAIM:

Sule

- 1. A method for transforming a plant comprising the steps of:
- (a) contacting the meristematic tissue of the plant with a medium comprising DNA;
 - (b) contacting an area of the plant below the megistematic tissue of step a) with a positive lead of a power source;
 - (c) contacting the medium comprising DNA with a negative lead of the power source; and
- 10 (d) applying a low amperage current from the power source, thereby causing the DNA to migrate from the medium to the cells of the meristematic tissue of the plant.
 - 2. The method of Claim 1, wherein the plant is a dicot.
 - 3. The method of Claim 2, wherein the plant is a soybean plant.
 - 4. The method of Claim 1, wherein/the plant is a monocot.
- 5. The method of Claim 1, wherein the plant is a seedling.
 - 6. The method of Claim 1, wherein the DNA is a plasmid vector.
 - 7. The method of Claim 6, wherein the plasmid vector is linearized.
 - 8. The method of Claim 6, wherein the plasmid contains the gene for barley oxalic acid oxidase.
 - 9. The method of Claim 1, wherein the current is about 0.01 to about 1.0 mA.
 - 10. The method of Claim 1, wherein the current is about 0.1 to about 0.5 mA.
- 35 11. The method of Claim 1, wherein the meristematic tissue is an apical meristem.

15

25

30

- 12. The method of Claim 1, wherein the meristematic tissue is a lateral meristem.
- 13. The method of Claim 1, wherein the meristematic tissue is a 5 meristematic dome.
 - 14. The method of Claim 1, wherein the area of the plant below the meristematic tissue is a root.
- 10 15. The method of Claim 1, wherein the area of the plant below the meristematic tissue is a stem.
 - 16. A trapsgenic plant produced by the method of Claim 1.

ADD >